

The LGMA recommends pre-harvest testing of product grown on fields where the following elevated risk factors are present:

Hazard	Current Required Minimum Distance	Elevated Risk Factors <i>If any of the following are present, test product before harvesting</i>
<b>Concentrated Animal Feeding Operation (CAFO)</b> 80,000+ animals	1 Mile	<ul style="list-style-type: none"> <li>- Distance &lt; 2 miles</li> <li>- Opportunity for water runoff through or from CAFO to irrigation water sources</li> <li>- Prevailing winds or other weather events that could potentially carry contaminants to fields</li> <li>- Filth fly infestation</li> </ul>
<b>CAFO</b> 1,000 - 80,000 animals	1200 Feet	<ul style="list-style-type: none"> <li>- Distance &lt; 1 mile</li> <li>- Opportunity for water runoff through or from CAFO to irrigation water sources</li> <li>- Opportunity for soil leaching</li> <li>- Prevailing winds and other weather events that could create potential contamination</li> <li>- Filth fly infestation</li> </ul>
<b>Animal Feed Operation (AFO)</b> < 1,000 animals	400 Feet (If composting on AFO site)  30 Feet (if no composting on AFO site)	<ul style="list-style-type: none"> <li>- Distance &lt; 1200 feet</li> <li>- Opportunity for water runoff through or from AFO to irrigation water sources</li> <li>- Opportunity for soil leaching</li> <li>- Prevailing winds or other weather events that could potentially carry contaminants to fields</li> <li>- Filth fly infestation</li> </ul>
<b>Grazing Lands</b>	30 Feet	<ul style="list-style-type: none"> <li>- Distance &lt; 1200 feet</li> <li>- Opportunity for water runoff through or from Grazing Land to irrigation water sources</li> <li>- Opportunity for soil leaching</li> <li>- Prevailing winds and other weather events that could create potential contamination</li> <li>- Filth fly infestation</li> </ul>
<b>Domestic / Hobby Farms</b> Includes non-commercial livestock and homes with hobby farms	30 Feet	<ul style="list-style-type: none"> <li>- Distance &lt; 800 feet</li> <li>- Opportunity for water runoff through or from Domestic/Hobby Farm to irrigation water sources</li> <li>- Opportunity for soil leaching</li> <li>- Prevailing winds or other weather events that could potentially carry contaminants to fields</li> <li>- Filth fly infestation</li> </ul>